





Q32
































oc or od is to be inclined at 30 degrees
 but neither o'c' nor o'd' is parallel in diag 3
 thus we need LOCUS
 but no where we have the TL
 since oa is // to xy, o'a' is TL of OA
 and that itself is TL of oc or od
 (since in real world OA=OB=OC=OD=...)




oc or od is to be inclined at 30 degrees
 but neither $o'c'$ nor $o'd'$ is parallel in diag 3
 thus we need LOCUS
 but no where we have the TL
 since oa is \parallel to xy , $o'a'$ is TL of OA
 and that itself is TL of oc or od
 (since in real world $OA=OB=OC=OD=...$)




oc or od is to be inclined at 30 degrees
 but neither o'c' nor o'd' is parallel in diag 3
 thus we need LOCUS
 but no where we have the TL
 since oa is // to xy , o'a' is TL of OA
 and that itself is TL of oc or od
 (since in real world OA=OB=OC=OD=...)




oc or od is to be inclined at 30 degrees
but neither o'c' nor o'd' is parallel in diag 3
thus we need LOCUS
but no where we have the TL
since oa is // to xy , o'a' is TL of OA
and that itself is TL of oc or od
(since in real world OA=OB=OC=OD=...)




oc or od is to be inclined at 30 degrees
 but neither o'c' nor o'd' is parallel in diag 3
 thus we need LOCUS
 but no where we have the TL
 since oa is // to xy, o'a' is TL of OA
 and that itself is TL of oc or od
 (since in real world OA=OB=OC=OD=...)




oc or od is to be inclined at 30 degrees
but neither o'c' nor o'd' is parallel in diag 3
thus we need LOCUS
but no where we have the TL
since oa is // to xy, o'a' is TL of OA
and that itself is TL of oc or od
(since in real world OA=OB=OC=OD=...)




oc or od is to be inclined at 30 degrees
but neither o'c' nor o'd' is parallel in diag 3
thus we need LOCUS
but no where we have the TL
since oa is // to xy, o'a' is TL of OA
and that itself is TL of oc or od
(since in real world OA=OB=OC=OD=...)



oc or od is to be inclined at 30 degrees
but neither o'c' nor o'd' is parallel in diag 3
thus we need LOCUS
but no where we have the TL
since oa is // to xy , o'a' is TL of OA
and that itself is TL of oc or od
(since in real world OA=OB=OC=OD=...)





oc or od is to be inclined at 30 degrees
but neither o'c' nor o'd' is parallel in diag 3
thus we need LOCUS
but no where we have the TL
since oa is // to xy, o'a' is TL of OA
and that itself is TL of oc or od
(since in real world OA=OB=OC=OD=...)



oc or od is to be inclined at 30 degrees
 but neither $o'c'$ nor $o'd'$ is parallel in diag 3
 thus we need LOCUS
 but no where we have the TL
 since oa is \parallel to xy , $o'a'$ is TL of OA
 and that itself is TL of oc or od
 (since in real world $OA=OB=OC=OD=...$)








oc or od is to be inclined at 30 degrees
 but neither o'c' nor o'd' is parallel in diag 3
 thus we need LOCUS
 but no where we have the TL
 since oa is // to xy, o'a' is TL of OA
 and that itself is TL of oc or od
 (since in real world OA=OB=OC=OD=...)





oc or od is to be inclined at 30 degrees
but neither o'c' nor o'd' is parallel in diag 3
thus we need LOCUS
but no where we have the TL
since oa is // to xy , o'a' is TL of OA
and that itself is TL of oc or od
(since in real world OA=OB=OC=OD=...)